




UNIVERSITÀ  
DI TRENTO



# Identifiability of centered Gaussian Mixtures and sums of powers of quadratics

**Alessandro Oneto**

with A. Taveira Blomenhofer, A. Casarotti, M. Michałek


arXiv:2204.09356

Joint Mathematics Meeting, Boston, January 6th

# Pearson's crabs: algebraic statistics in 1893

(1983)

K. Pearson wanted to explain an asymmetry observed in data about crabs in Naples'





III. *Contributions to the Mathematical Theory of Evolution.*  
By KARL PEARSON, *University College, London.*  
*Communicated by Professor HENRICI, F.R.S.*  
Received October 18,—Read November 16, 1893.

normal shape, and it becomes important to determine the direction and amount of such deviation. The asymmetry may arise from the fact that the units grouped together in the measured material are not really homogeneous. It may happen that we have a mixture of 2, 3, . . .  $n$  homogeneous groups, each of which deviates about its own mean symmetrically and in a manner represented with sufficient accuracy by the normal curve. Thus an abnormal frequency-curve may be really built up of normal curves having parallel but not necessarily coincident axes and different parameters.

# Pearson's crabs: algebraic statistics in 1893

(1983)

K. Pearson wanted to explain an asymmetry observed in data about crabs in Naples'



# Pearson's crabs: algebraic statistics in 1894

(1984)

K. Pearson wanted to explain an asymmetry observed in data about crabs in Naples'

## Method of moments.

$$z_1 + z_2 = 1 \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad (8).$$

$$\gamma_1 z_1 + \gamma_2 z_2 = 0 \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad (9).$$

$$\gamma_1^2 z_1 (1 + u_1^2) + \gamma_2^2 z_2 (1 + u_2^2) = \mu_2 \quad . \quad . \quad . \quad . \quad . \quad (10).$$

$$\gamma_1^3 z_1 (1 + 3u_1^2) + \gamma_2^3 z_2 (1 + 3u_2^2) = \mu_3 \cdot \cdot \cdot \cdot (11).$$

$$\gamma_1^4 z_1 (1 + 6u_1^2 + 3u_1^4) + \gamma_2^4 z_2 (1 + 6u_2^2 + 3u_2^4) = \mu_4 \quad . \quad . \quad (12).$$

$$\gamma_1^5 z_1 (1 + 10u_1^2 + 15u_1^4) + \gamma_2^5 z_2 (1 + 10u_2^2 + 15u_2^4) = \mu_5 \quad . \quad (13).$$

Equations (8)-(13) give the complete solution of the problem.\* After several trials, I find that the elimination of  $z_1, z_2, u_1, u_2$  from these equations, and the determination of equations giving  $\gamma_1\gamma_2$  and  $\gamma_1 + \gamma_2$  appear to lead to a resulting equation of the lowest possible order.




Figure 4.2.1.: Approximation of crabs data by a mixture of two Gaussians.

# Moments

Consider a random variable  $X = (X_1, \dots, X_n)$  defined by a density function  $f(X) : \mathbb{R}^n \rightarrow \mathbb{R}$ .

For  $\mathbf{i} = (i_1, \dots, i_n) \in \mathbb{N}^n$ , the **i-moment** of  $X$  is:

$$m_{\mathbf{i}} = \mathbb{E}[X_1^{i_1} \cdots X_n^{i_n}] = \int_{\mathbb{R}^n} x^{\mathbf{i}} f(x) dx.$$

For  $d \in \mathbb{N}$ , the **d-moment** of  $X$  is the degree- $d$  polynomial:

$$m_d = \mathbb{E}[\langle X, Y \rangle^d] = \sum_{\mathbf{i} \in \mathbb{N}^n, i_1 + \dots + i_n = d} \frac{d!}{i_1! \cdots i_n!} m_{\mathbf{i}} Y^{\mathbf{i}} \in \mathbb{R}[Y].$$

# Moments

Consider a random variable  $X = (X_1, \dots, X_n)$  defined by a density function  $f(X) : \mathbb{R}^n \rightarrow \mathbb{R}$ .

For  $\mathbf{i} = (i_1, \dots, i_n) \in \mathbb{N}^n$ , the **i-moment** of  $X$  is:

$$m_{\mathbf{i}} = \mathbb{E}[X_1^{i_1} \cdots X_n^{i_n}] = \int_{\mathbb{R}^n} x^{\mathbf{i}} f(x) dx.$$

For  $d \in \mathbb{N}$ , the **d-moment** of  $X$  is the degree- $d$  polynomial:

$$m_d = \mathbb{E}[\langle X, Y \rangle^d] = \sum_{\mathbf{i} \in \mathbb{N}^n, i_1 + \dots + i_n = d} \frac{d!}{i_1! \cdots i_n!} m_{\mathbf{i}} Y^{\mathbf{i}} \in \mathbb{R}[Y].$$

A compact way to look at moments is via the **moment generating function**:


$$\mathbb{E}[\exp(\langle X, Y \rangle)] = \sum_d \frac{1}{d!} \mathbb{E}[\langle X, Y \rangle^d] = \sum_d \frac{1}{d!} m_d \in \mathbb{R}[[Y]].$$

# Moments of Gaussian distributions

Consider a multivariate Gaussian distribution  $X \sim \mathcal{N}(\mu, \Sigma)$ , i.e.,

$$f(x) = \frac{1}{\sqrt{\det(2\pi\Sigma)}} \exp\left(-\frac{1}{2}(x - \mu)^T \Sigma^{-1}(x - \mu)\right).$$

- $\mu = (\mu^1, \dots, \mu^n) \in \mathbb{R}^n$ : **mean vector**;
- $\Sigma = (\sigma_{ij})_{i,j} \in \mathbb{R}^{n \times n}$ : **covariance matrix**, i.e., symmetric positive semidefinite matrix.



WikiMedia Commons

# Moments of Gaussian distributions

Consider a multivariate Gaussian distribution  $X \sim \mathcal{N}(\mu, \Sigma)$ , i.e.,

$$f(x) = \frac{1}{\sqrt{\det(2\pi\Sigma)}} \exp \left( -\frac{1}{2}(x - \mu)^T \Sigma^{-1}(x - \mu) \right).$$

- $\mu = (\mu^1, \dots, \mu^n) \in \mathbb{R}^n$ : **mean vector**;
- $\Sigma = (\sigma_{ij})_{i,j} \in \mathbb{R}^{n \times n}$ : **covariance matrix**, i.e., symmetric positive semidefinite matrix.


They define the homogeneous polynomials:

$$\ell = \mu^T X = \mu^1 Y_1 + \dots + \mu^n Y_n;$$

$$q = X^T \Sigma X = \sum_{ij} \sigma_{ij} X_i X_j.$$

Then, the **moment generating function** is

$$\mathbb{E}[\exp(\langle X, Y \rangle)] = \sum_d \frac{1}{d!} m_d = \exp \left( \ell + \frac{1}{2} q \right) = \sum_d \frac{1}{d!} \left( \ell + \frac{1}{2} q \right)^d.$$




WikiMedia Commons

# Moments of Gaussian distributions

Consider a multivariate Gaussian distribution  $X \sim \mathcal{N}(\mu, \Sigma)$ , i.e.,

$$f(x) = \frac{1}{\sqrt{\det(2\pi\Sigma)}} \exp \left( -\frac{1}{2}(x - \mu)^T \Sigma^{-1}(x - \mu) \right).$$



- $\mu = (\mu^1, \dots, \mu^n) \in \mathbb{R}^n$ : **mean vector**;
- $\Sigma = (\sigma_{ij})_{i,j} \in \mathbb{R}^{n \times n}$ : **covariance matrix**, i.e., symmetric

They define the homogeneous polynomials:

$$\ell = \mu^T X = \mu^1 Y_1 + \dots + \mu^n Y_n;$$

| $d$ | degree- $d$ moment, $m_d$                    |
|-----|----------------------------------------------|
| 1   | $\ell$                                       |
| 2   | $\ell^2 + q$                                 |
| 3   | $\ell^3 + 3\ell q$                           |
| 4   | $\ell^4 + 6\ell^2 q + 3q^2$                  |
| 5   | $\ell^5 + 10\ell^3 q + 15q^2 \ell$           |
| 6   | $\ell^6 + 15\ell^4 q + 45\ell^2 q^2 + 15q^4$ |

Then, the **moment generating function** is

$$\mathbb{E}[\exp(\langle X, Y \rangle)] = \sum_d \frac{1}{d!} m_d = \exp \left( \ell + \frac{1}{2} q \right) = \sum_d \frac{1}{d!} \left( \ell + \frac{1}{2} q \right)^d.$$

# Moments of mixtures of Gaussian distributions

Consider a multivariate Gaussian distributions

$$X_1 \sim \mathcal{N}(\mu_1, \Sigma_1), \dots, X_m \sim \mathcal{N}(\mu_m, \Sigma_m),$$

and consider a **mixture**

$$X = \lambda_1 X_1 + \dots + \lambda_m X_m, \quad \text{where } \lambda_i \in (0,1), \quad \sum_i \lambda_i = 1.$$

Then, the moment generating function is

$$\begin{aligned} \mathbb{E}[\exp(\langle X, Y \rangle)] &= \sum_{i=1}^m \lambda_i \mathbb{E}[\exp(\langle X_i, Y \rangle)] \\ &= \exp(\lambda_1 \mu_1 + \dots + \lambda_m \mu_m) \exp\left(\frac{1}{2}(\lambda_1 q_1 + \dots + \lambda_m q_m)\right) = \sum_d \frac{1}{d!} \sum_{i=1}^m \lambda_i \left(\ell_i + \frac{1}{2} q_i\right)^d \end{aligned}$$

# Identifiability of mixtures of Gaussian distributions

Consider a multivariate Gaussian distributions


$$X_1 \sim \mathcal{N}(\mu_1, \Sigma_1), \dots, X_m \sim \mathcal{N}(\mu_m, \Sigma_m),$$

and consider a **mixture**

$$X = \lambda_1 X_1 + \dots + \lambda_m X_m, \quad \text{where } \lambda_i \in (0,1), \quad \sum_i \lambda_i = 1.$$

For some  $d \in \mathbb{N}$ , consider the **inverse problem** of the map sending the parameters of the model to finitely-many moments:

$$(\lambda_1, \dots, \lambda_m, \ell_1, \dots, \ell_m, q_1, \dots, q_m) \mapsto (m_1, m_2, \dots, m_d).$$



# Identifiability of mixtures of Gaussian distributions

Consider a multivariate Gaussian distributions

$$X_1 \sim \mathcal{N}(\mu_1, \Sigma_1), \dots, X_m \sim \mathcal{N}(\mu_m, \Sigma_m),$$


and consider a **mixture**

$$X = \lambda_1 X_1 + \dots + \lambda_m X_m, \quad \text{where } \lambda_i \in (0,1), \quad \sum_i \lambda_i = 1.$$

For some  $d \in \mathbb{N}$ , consider the **inverse problem** of the map sending the parameters of the model to finitely-many moments:

$$(\lambda_1, \dots, \lambda_m, \ell_1, \dots, \ell_m, q_1, \dots, q_m) \mapsto (m_1, m_2, \dots, m_d).$$

**Question.** Given finitely-many moments  $(m_1, m_2, \dots, m_d)$ , are the parameters of the mixture model *generically* identified (up to permutation)?



# Identifiability of mixtures of Gaussian distributions

## the case with identical covariance

Consider a multivariate Gaussian distributions


$$X_1 \sim \mathcal{N}(\mu_1, \Sigma), \dots, X_m \sim \mathcal{N}(\mu_m, \Sigma),$$

and consider a **mixture**

$$X = \lambda_1 X_1 + \dots + \lambda_m X_m, \quad \text{where } \lambda_i \in (0,1), \quad \sum_i \lambda_i = 1.$$

If  $\ell_i = x^T \mu_i$  and  $q = x^T \Sigma x$ , then we have that the first moments are

$$m_1 = \sum_{i=1}^m \ell_i, \quad m_2 = \sum_{i=1}^m \ell_i^2 + mq, \quad m_3 = \sum_{i=1}^m \ell_i^3 + 3mq \sum_{i=1}^m \ell_i.$$



# Identifiability of mixtures of Gaussian distributions

## the case with identical covariance

Consider a multivariate Gaussian distributions

$$X_1 \sim \mathcal{N}(\mu_1, \Sigma), \dots, X_m \sim \mathcal{N}(\mu_m, \Sigma),$$


and consider a **mixture**

$$X = \lambda_1 X_1 + \dots + \lambda_m X_m, \quad \text{where } \lambda_i \in (0,1), \quad \sum_i \lambda_i = 1.$$

If  $\ell_i = x^T \mu_i$  and  $q = x^T \Sigma x$ , then we have that the first moments are

$$m_1 = \sum_{i=1}^m \ell_i, \quad m_2 = \sum_{i=1}^m \ell_i^2 + m q, \quad m_3 = \sum_{i=1}^m \ell_i^3 + 3 m q \sum_{i=1}^m \ell_i.$$

By shifting the space, we may assume  $\sum_{i=1}^m \ell_i = 0$ .



# Identifiability of mixtures of Gaussian distributions

## the case with identical covariance

Consider a multivariate Gaussian distributions

$$X_1 \sim \mathcal{N}(\mu_1, \Sigma), \dots, X_m \sim \mathcal{N}(\mu_m, \Sigma),$$

and consider a **mixture**


$$X = \lambda_1 X_1 + \dots + \lambda_m X_m, \quad \text{where } \lambda_i \in (0,1), \quad \sum_i \lambda_i = 1.$$

If  $\ell_i = x^T \mu_i$  and  $q = x^T \Sigma x$ , then we have that the first moments are

$$m_1 = \sum_{i=1}^m \ell_i, \quad m_2 = \sum_{i=1}^m \ell_i^2 + m q, \quad m_3 = \sum_{i=1}^m \ell_i^3 + \cancel{3 m \sum_{i=1}^m \ell_i q}.$$

By shifting the space, we may assume  $\sum_{i=1}^m \ell_i = 0$ .

**Question.** Is  $m_3 = \sum_{i=1}^m \ell_i^3$  generically identifiable? I.e., given a general cubic homogeneous polynomial, is its expression as sum of cubes unique up to permutation?



# Identifiability of mixtures of Gaussian distributions

## the case with identical covariance

**Question.** Is the map  $m_3 = \sum_{i=1}^m \ell_i^3$  generically identifiable? I.e., given a general cubic homogeneous polynomial, is its expression as sum of cubes unique up to permutation?

Given a homogeneous polynomial  $f \in \text{Sym}^d(\mathbb{C}^n)$ , its **Waring rank** is

$$\text{rk}(f) = \min_m \left\{ f = \sum_{i=1}^m \ell_i^d : \ell_i \in \text{Sym}^1(\mathbb{C}^n) \right\}.$$

**Theorem (Chiantini-Ottaviani-Vannieuwenhoven, 2017).** For  $d \geq 3$ , the general homogeneous polynomial  $f \in \text{Sym}^d(\mathbb{C}^n)$  of Waring rank smaller than the generic one has a unique minimal Waring decomposition except for  $(d, n, m) \in \{(6, 2, 9), (4, 3, 8), (3, 5, 9)\}$ .

# Identifiability of mixtures of Gaussian distributions

## the centered case

Consider a multivariate Gaussian distributions


$$X_1 \sim \mathcal{N}(0, \Sigma_1), \dots, X_m \sim \mathcal{N}(0, \Sigma_m),$$

and consider a **mixture**

$$X = \lambda_1 X_1 + \dots + \lambda_m X_m, \quad \text{where } \lambda_i \in (0,1), \quad \sum_i \lambda_i = 1.$$

Since  $\ell_1 = \dots = \ell_m = 0$  and  $q = x^T \Sigma x$ , then we have that the moments are

$$m_i = \begin{cases} 0 & \text{for } i = 2d + 1 \\ q_1^d + \dots + q_m^d & \text{for } i = 2d \end{cases}.$$



| $d$ | degree- $d$ moment, $m_d$                    |
|-----|----------------------------------------------|
| 1   | $\ell$                                       |
| 2   | $\ell^2 + q$                                 |
| 3   | $\ell^3 + 3\ell q$                           |
| 4   | $\ell^4 + 6\ell^2 q + 3q^2$                  |
| 5   | $\ell^5 + 10\ell^3 q + 15q^2 \ell$           |
| 6   | $\ell^6 + 15\ell^4 q + 45\ell^2 q^2 + 15q^4$ |

# Identifiability of mixtures of Gaussian distributions

## the centered case

Consider a multivariate Gaussian distributions

$$X_1 \sim \mathcal{N}(0, \Sigma_1), \dots, X_m \sim \mathcal{N}(0, \Sigma_m),$$


and consider a **mixture**

$$X = \lambda_1 X_1 + \dots + \lambda_m X_m, \quad \text{where } \lambda_i \in (0,1), \quad \sum_i \lambda_i = 1.$$

Since  $\ell_1 = \dots = \ell_m = 0$  and  $q = x^T \Sigma x$ , then we have that the moments are

$$m_i = \begin{cases} 0 & \text{for } i = 2d + 1 \\ q_1^d + \dots + q_m^d & \text{for } i = 2d \end{cases}.$$

**Question.** Is  $m_{2d} = \sum_{i=1}^m q_i^d$  generically identifiable?  
I.e., given a general homogeneous polynomial of even degree, is its expression as sum of powers of quadrics unique up to permutation?



# Identifiability of mixtures of Gaussian distributions

## the centered case

**Question.** Is  $m_{2d} = \sum_{i=1}^m q_i^d$  generically identifiable?  
I.e., given a general homogeneous polynomial of even degree, is its expression as sum of powers of quadrics unique up to permutation?

**Observation.** For degree-2 and degree-4 moments we cannot have identifiability. Indeed:

- $q_1 + q_2 = (q_1 + q') + (q_2 - q')$ , for any quadric  $q'$ ;
- $q_1^2 + q_2^2 = \frac{1}{2}(q_1 + q_2)^2 + \frac{1}{2}(q_1 - q_2)^2$ .

# Identifiability of mixtures of Gaussian distributions

## the centered case

**T**heorem (Taveira Blomenhofer-Casarotti-Michałek-Oneto, 2022\*). For  $m, n \in \mathbb{N}$  such that

$$m \leq \min \left\{ \frac{\binom{n+5}{6}}{\binom{n+1}{2}} - \binom{n+1}{2} - 1, \binom{n+1}{2} + 1 \right\}$$

the general sextic  $f = \sum_{i=1}^m q_i^3$  is identifiable.

# Identifiability of mixtures of Gaussian distributions

## the centered case

**T**heorem (Taveira Blomenhofer-Casarotti-Michalek-Oneto, 2022\*). For  $m, n \in \mathbb{N}$  such that

$$m \leq \min \left\{ \frac{\binom{n+5}{6}}{\binom{n+1}{2}} - \binom{n+1}{2} - 1, \binom{n+1}{2} + 1 \right\}$$

the general sextic  $f = \sum_{i=1}^m q_i^3$  is identifiable.

**C**orollary (Taveira Blomenhofer-Casarotti-Michalek-Oneto, 2022\*).  
For  $m, n \in \mathbb{N}$  such that the Theorem holds then:

1. the uniformly weighted Gaussian mixture  $X = \frac{1}{m}(X_1 + \dots + X_m)$  is generically identified by its degree-6 moment;
2. the general Gaussian mixture  $X = \lambda_1 X_1 + \dots + \lambda_m X_m$  is generically identified by its degree-6 and degree-4 moments.

# Identifiability of mixtures of Gaussian distributions


## the centered case

**T**heorem (Taveira Blomenhofer-Casarotti-Michalek-Oneto, 2022\*). For  $m, n \in \mathbb{N}$  such that

$$m \leq \min \left\{ \frac{\binom{n+5}{6}}{\binom{n+1}{2}} - \binom{n+1}{2} - 1, \binom{n+1}{2} + 1 \right\}$$

the general sextic  $f = \sum_{i=1}^m q_i^3$  is identifiable.

$$\frac{\binom{n+5}{6}}{\binom{n+1}{2}}$$



# From non-defectivity to identifiability

Given any algebraic variety  $X \subset \mathbb{P}^N$ , we define:

- For any point  $q \in \mathbb{P}^N$ , the  **$X$ -rank** of  $q$  is  $\text{rk}_X(q) = \min_m \{q \in \langle p_1, \dots, p_m \rangle : p_i \in X\}$ .
- The  **$m$ -secant variety** of  $X$  is  $\sigma_m(X) = \overline{\bigcup_{p_1, \dots, p_m \in X} \langle p_1, \dots, p_m \rangle} = \overline{\{q : \text{rk}_X(q) \leq m\}}$ .

# From non-defectivity to identifiability

Given any algebraic variety  $X \subset \mathbb{P}^N$ , we define:

- For any point  $q \in \mathbb{P}^N$ , the  **$X$ -rank** of  $q$  is  $\text{rk}_X(q) = \min_m \{q \in \langle p_1, \dots, p_m \rangle : p_i \in X\}$ .
- The  **$m$ -secant variety** of  $X$  is  $\sigma_m(X) = \overline{\bigcup_{p_1, \dots, p_m \in X} \langle p_1, \dots, p_m \rangle} = \overline{\{q : \text{rk}_X(q) \leq m\}}$ .

The variety  $X$  is said  **$m$ -defective** if  $\dim \sigma_m(X) < \min\{N, m \dim(X) + m - 1\}$ .

The variety  $X$  is said  **$m$ -identifiable** if the general element  $q \in \sigma_m(X)$  is identifiable.

# From non-defectivity to identifiability

Given any algebraic variety  $X \subset \mathbb{P}^N$ , we define:

- For any point  $q \in \mathbb{P}^N$ , the  **$X$ -rank** of  $q$  is  $\text{rk}_X(q) = \min_m \{q \in \langle p_1, \dots, p_m \rangle : p_i \in X\}$ .
- The  **$m$ -secant variety** of  $X$  is  $\sigma_m(X) = \overline{\bigcup_{p_1, \dots, p_m \in X} \langle p_1, \dots, p_m \rangle} = \overline{\{q : \text{rk}_X(q) \leq m\}}$ .

The variety  $X$  is said  **$m$ -defective** if  $\dim \sigma_m(X) < \min\{N, m \dim(X) + m - 1\}$ .

The variety  $X$  is said  **$m$ -identifiable** if the general element  $q \in \sigma_m(X)$  is identifiable.

**Theorem (Casarotti-Mella, 2020).** If  $X$  is smooth,  $m > 2 \dim(X)$ ,  $m \dim(X) + m - 1 < N$ , then non  $m$ -defectivity implies  $(m - 1)$ -identifiability.

# From non-defectivity to identifiability

Given any algebraic variety  $X \subset \mathbb{P}^N$ , we define:

- For any point  $q \in \mathbb{P}^N$ , the  **$X$ -rank** of  $q$  is  $\text{rk}_X(q) = \min_m \{q \in \langle p_1, \dots, p_m \rangle : p_i \in X\}$ .
- The  **$m$ -secant variety** of  $X$  is  $\sigma_m(X) = \overline{\bigcup_{p_1, \dots, p_m \in X} \langle p_1, \dots, p_m \rangle} = \overline{\{q : \text{rk}_X(q) \leq m\}}$ .

The variety  $X$  is said  **$m$ -defective** if  $\dim \sigma_m(X) < \min\{N, m \dim(X) + m - 1\}$ .

The variety  $X$  is said  **$m$ -identifiable** if the general element  $q \in \sigma_m(X)$  is identifiable.

**Theorem (Massarenti-Mella, 2022).** If  $X$  is smooth,  ~~$m > 2 \dim(X)$~~ ,  $m \dim(X) + m - 1 < N$ , then non  $m$ -defectivity implies  $(m - 1)$ -identifiability.

# Identifiability of mixtures of Gaussian distributions

## the centered case

**T**heorem (Taveira Blomenhofer-Casarotti-Michalek-Oneto, 2022\*). For  $m, n \in \mathbb{N}$  such that

$$m \leq \min \left\{ \frac{\binom{n+5}{6}}{\binom{n+1}{2}} - \binom{n+1}{2} - 1, \binom{n+1}{2} + 1 \right\}$$

the general sextic  $f = \sum_{i=1}^m q_i^3$  is identifiable.


from non-defectivity  
to identifiability

$$\frac{\binom{n+5}{6}}{\binom{n+1}{2}}$$

the expected number of cubes needed by the general sextic

tangential contact locus

Massarenti-Mella, 2022



# Non-defectivity of sums of powers


## and Fröberg's Conjecture

Let  $V_{d,k,n} = \{[g^k] : g \in \mathbb{P}\mathrm{Sym}^d(\mathbb{C}^n)\} \subset \mathbb{P}\mathrm{Sym}^{dk}(\mathbb{C}^n)$

.

**C**onjecture (Ottaviani, 2017).

For  $k \geq 3$ ,  $d \geq 2$ , the variety  $V_{d,k,n}$  is never  $m$ -defective.



# Non-defectivity of sums of powers


## and Fröberg's Conjecture

Let  $V_{d,k,n} = \{[g^k] : g \in \mathbb{P}\mathrm{Sym}^d(\mathbb{C}^n)\} \subset \mathbb{P}\mathrm{Sym}^{dk}(\mathbb{C}^n)$

.

**C**onjecture (Ottaviani, 2017).

For  $k \geq 3$ ,  $d \geq 2$ , the variety  $V_{d,k,n}$  is never  $m$ -defective.



**L**emma (Terracini, 1911). Let  $X \subset \mathbb{P}^N$  smooth.

Given general points  $p_1, \dots, p_m \in X$  and general  $q \in \langle p_1, \dots, p_m \rangle$ , then

$$T_q \sigma_m(X) = \langle T_{p_1}(X), \dots, T_{p_m}(X) \rangle.$$

# Non-defectivity of sums of powers


## and Fröberg's Conjecture

Let  $V_{d,k,n} = \{[g^k] : g \in \mathbb{P}\mathrm{Sym}^d(\mathbb{C}^n)\} \subset \mathbb{P}\mathrm{Sym}^{dk}(\mathbb{C}^n)$

.

**C**onjecture (Ottaviani, 2017).

For  $k \geq 3$ ,  $d \geq 2$ , the variety  $V_{d,k,n}$  is never  $m$ -defective.



Note that  $\frac{d}{dt} \big|_{t=0} (g + th)^k = kg^{k-1}h$ .

Given general forms  $g_1, \dots, g_m \in \mathrm{Sym}^d(\mathbb{C}^n)$  and  $I = (g_1^{k-1}, \dots, g_m^{k-1})$ , the **general tangent space** to  $\sigma_m(V_{d,k,n})$  is

$$\mathbb{P}(I \cap \mathrm{Sym}^{dk}(\mathbb{C}^n)) = \mathbb{P}[(g_1^{k-1}, \dots, g_m^{k-1})]_{kd}$$

# Non-defectivity of sums of powers

## and Fröberg's Conjecture

**C**onjecture (Nicklasson, 2017).

Given general forms  $g_1, \dots, g_m \in \text{Sym}^d(\mathbb{C}^n)$  and  $I = (g_1^{k-1}, \dots, g_m^{k-1})$ ,

$$\sum_i \dim (I \cap \text{Sym}^i(\mathbb{C}^n)) t^i = \left[ \frac{(1 - t^{d(k-1)})^m}{(1 - t)^n} \right],$$

where  $[ \cdot ]$  means to truncate the power series before the first non-positive coefficient.  
I.e., the ideal  $I$  has the minimal Hilbert function prescribed by Fröberg's Conjecture.

R. Fröberg. *An inequality for Hilbert series of graded algebras*. Mathematica Scandinavica. 1985

A. Oneto, *Waring-type problems for polynomials*. PhD thesis, Stockholm U., 2016

L. Nicklasson. *On the Hilbert series of ideals generated by generic forms*. Communications in Algebra. 2017

A note on Fröberg's conjecture for forms of equal degrees. *Comptes Rendus Mathematique*. 2017

# Non-defectivity of sums of powers

## and Fröberg's Conjecture

**C**onjecture (Nicklasson, 2017).

Given general forms  $g_1, \dots, g_m \in \text{Sym}^d(\mathbb{C}^n)$  and  $I = (g_1^{k-1}, \dots, g_m^{k-1})$ ,

$$\sum_i \dim(I \cap \text{Sym}^i(\mathbb{C}^n)) t^i = \left[ \frac{(1 - t^{d(k-1)})^m}{(1 - t)^n} \right],$$

where  $[\cdot]$  means to truncate the power series before the first non-positive coefficient.  
I.e., the ideal  $I$  has the minimal Hilbert function prescribed by Fröberg's Conjecture.

$$\text{coeff}_{kd} \frac{(1 - t^{d(k-1)})^m}{(1 - t)^n} = \binom{n + dk - 1}{dk} - m \binom{n + d - 1}{d} = \exp . \text{codim} \sigma_m(V_{d,k,n})$$

R. Fröberg. *An inequality for Hilbert series of graded algebras*. Mathematica Scandinavica. 1985

A. Oneto, *Waring-type problems for polynomials*. PhD thesis, Stockholm U., 2016

L. Nicklasson. *On the Hilbert series of ideals generated by generic forms*. Communications in Algebra. 2017

A note on Fröberg's conjecture for forms of equal degrees. *Comptes Rendus Mathématique*. 2017

# Non-defectivity of sums of powers

## and Fröberg's Conjecture

**C**onjecture (Nicklasson, 2017).

Given general forms  $g_1, \dots, g_m \in \text{Sym}^d(\mathbb{C}^n)$  and  $I = (g_1^{k-1}, \dots, g_m^{k-1})$ ,

$$\sum_i \dim (I \cap \text{Sym}^i(\mathbb{C}^n)) t^i = \left[ \frac{(1 - t^{d(k-1)})^m}{(1 - t)^n} \right],$$

where  $[ \cdot ]$  means to truncate the power series before the first non-positive coefficient.  
I.e., the ideal  $I$  has the minimal Hilbert function prescribed by Fröberg's Conjecture.

**T**heorem (Nenashev, 2015). If  $m \leq \frac{\dim \text{Sym}^{d(k-1)}(\mathbb{C}^n)}{\dim \text{Sym}^d(\mathbb{C}^n)} - \dim \text{Sym}^d(\mathbb{C}^n)$ ,  
then the Conjecture holds.

R. Fröberg. *An inequality for Hilbert series of graded algebras*. Mathematica Scandinavica. 1985


A. Oneto, *Waring-type problems for polynomials*. PhD thesis, Stockholm U., 2016

L. Nicklasson. *On the Hilbert series of ideals generated by generic forms*. Communications in Algebra. 2017

A note on Fröberg's conjecture for forms of equal degrees. *Comptes Rendus Mathématique*. 2017



UNIVERSITÀ  
DI TRENTO



# Thank you!

---

A. Taveira Blomenhofer, A. Casarotti, M. Michałek, **A. Oneto**.

*Identifiability for mixtures of centered Gaussians and sums of powers of quadratics.*

arXiv:2204.09356v2